

OWUAM-029

July 10, 1998

DRINFELD TWIST FOR TWO-PARAMETRIC DEFORMATION OF $gl(2)$ AND $sl(1/2)$ [†]

N. Aizawa

Department of Applied Mathematics, Osaka Women's University
Sakai, Osaka 590-0035, Japan

Abstract

Drinfeld twist is applied to the Lie algebra $gl(2)$ so that a two-parametric deformation of it is obtained, which is identical to the Jordanian deformation of the $gl(2)$ obtained by Aneva *et al.* The same twist element is applied to deform the Lie superalgebra $sl(1/2)$, since the $gl(2)$ is embedded into the $sl(1/2)$. By making use of the FRT-formalism, we construct a deformation of the Lie supergroup $SL(1/2)$.

[†]Presented at the 7th International Colloquium "Quantum Groups and Integrable Systems", Prague, 18–20 June 1998.

1 Introduction

It is well known that deformation of Lie groups or Lie algebras is not unique. As for $gl(2)$, its multiparametric quantum deformation is classified recently [1]. In this note, a two-parametric deformation of $gl(2)$ by the Drinfeld twist [2] is considered and it is shown that the deformed $gl(2)$ is identical to the one obtained before [4]. The twist for $gl(2)$ used here is also applicable to deform the Lie superalgebra $sl(1/2)$, since $gl(2) \subset sl(1/2)$. The universal R-matrix for the deformed $sl(1/2)$ is constructed according to the method of twisting. This enables us, using the FRT-formalism, to construct a deformation of the Lie supergroup $SL(1/2)$. The same scenario has been carried out for a one-parametric deformation of $sl(2)$ and $osp(1/2)$ (Note that $sl(2) \subset osp(1/2)$)[3]. The present work follows the line adapted in [3].

The deformation by the Drinfeld twist has been developed in recent years. For example, multiparametric twists for the Drinfeld-Jimbo deformation of simple Lie algebras [8], one-parametric twist for $sl(2)$ [9], and twist for Poincaré algebra [10], Heisenberg algebra [11], esoteric quantum groups [12], $sl(N)$ [13], and $osp(1/2)$ [3, 14] have been considered.

2 Drinfeld twist

This section is devoted to a brief review of the Drinfeld twist. Drinfeld develops it in his study of quasi Hopf algebras, we however restrict ourselves to ordinary Hopf algebras throughout this note.

Let $\mathcal{A}$ be a Hopf algebra with coproduct Δ_0 , counit ϵ_0 and antipode S_0 . Let $\mathcal{F}$ be an invertible element in $\mathcal{A} \otimes \mathcal{A}$ satisfying the conditions

$$\begin{aligned} (\epsilon_0 \otimes id)(\mathcal{F}) &= (id \otimes \epsilon_0)(\mathcal{F}) = 1, \\ \mathcal{F}_{12}(\Delta_0 \otimes id)(\mathcal{F}) &= \mathcal{F}_{23}(id \otimes \Delta_0)(\mathcal{F}), \end{aligned} \tag{2.1}$$

then we obtain a new Hopf algebra $\mathcal{H}$ with the same algebraic relations as $\mathcal{A}$. However, Hopf algebraic mappings of $\mathcal{H}$ are different, they are twisted by the twist element $\mathcal{F}$. The coproduct, the counit and the antipode for $\mathcal{H}$ are given by

$$\Delta = \mathcal{F}\Delta_0\mathcal{F}^{-1}, \quad \epsilon = \epsilon_0, \quad S = uS_0u^{-1}, \tag{2.2}$$

where $u = m(id \otimes S_0)(\mathcal{F})$.

When the algebra $\mathcal{A}$ has a universal R-matrix $\mathcal{R}_0$, the universal R-matrix $\mathcal{R}$ for $\mathcal{H}$ is given by

$$\mathcal{R} = \mathcal{F}_{21}\mathcal{R}_0\mathcal{F}^{-1}. \tag{2.3}$$

This shows that if $\mathcal{A}$ is cocommutative, then $\mathcal{H}$ is a triangular Hopf algebra. This is the case that we would like to consider, since we shall start with a Lie algebra.

3 Twist for $gl(2)$

The Lie algebra $gl(2)$ has elements Z, H , and $X_{\pm}$ and defined by the relations

$$[H, X_{\pm}] = \pm 2X_{\pm}, \quad [X_+, X_-] = H, \quad [Z, \bullet] = 0. \quad (3.1)$$

A twist element for $gl(2)$ is given by

$$\mathcal{F} = \exp\left(\frac{g}{2h}\sigma \otimes Z\right) \exp\left(-\frac{1}{2}H \otimes \sigma\right), \quad (3.2)$$

with

$$\sigma \equiv -\ln(1 - 2hX_+),$$

where h and g are deformation parameters.

According to (2.2), the coproduct and the antipode are deformed and they read

$$\begin{aligned} \Delta(Z) &= Z \otimes 1 + 1 \otimes Z, \\ \Delta(H) &= H \otimes e^{\sigma} + 1 \otimes H + \frac{g}{h}(1 - e^{\sigma}) \otimes Ze^{\sigma}, \\ \Delta(X_+) &= X_+ \otimes 1 + e^{-\sigma} \otimes X_+, \\ \Delta(X_-) &= X_- \otimes e^{\sigma} + 1 \otimes X_- - hH \otimes e^{\sigma}H - \frac{h}{2}H(H+2) \otimes e^{\sigma}(e^{\sigma}-1) \\ &\quad + g(e^{\sigma}-1) \otimes Ze^{\sigma}H + g(H - e^{\sigma} + 1) \otimes Ze^{\sigma} \\ &\quad + g(e^{\sigma}-1)(H + e^{\sigma} + 1) \otimes Ze^{2\sigma} \\ &\quad - \frac{g^2}{2h}(e^{\sigma}-1) \otimes Z^2e^{\sigma} - \frac{g^2}{2h}(e^{\sigma}-1)^2 \otimes Z^2e^{2\sigma}, \\ S(Z) &= -Z, \\ S(H) &= -He^{-\sigma} + \frac{g}{h}Z(e^{-\sigma}-1), \\ S(X_+) &= -Xe^{\sigma}, \\ S(X_-) &= -\left\{X_- + \frac{h}{2}H^2(e^{-\sigma}+1) - hH(e^{-\sigma}-1) - gHZe^{-\sigma} \right. \\ &\quad \left. + gZ(e^{-\sigma}-1) + \frac{g^2}{2h}(e^{-\sigma}-1)Z^2\right\}e^{-\sigma}. \end{aligned} \quad (3.3)$$

The universal R-matrix for the non-cocommutative coproduct given above reads

$$\mathcal{R} = \exp\left(\frac{g}{2h}Z \otimes \sigma\right) \exp\left(-\frac{1}{2}\sigma \otimes H\right) \exp\left(\frac{1}{2}H \otimes \sigma\right) \exp\left(-\frac{g}{2h}\sigma \otimes Z\right). \quad (3.4)$$

A counit is necessary for a Hopf algebra, it is undeformed, that is, $\epsilon(Z) = \epsilon(H) = \epsilon(X_{\pm}) = 0$. Thus we arrive to the definition

Definition 1. The triangular Hopf algebra generated by $\{Z, H, X_{\pm}\}$ satisfying the relations (3.1) and (3.3) is said to be the two-parametric deformation of $\mathcal{U}(gl(2))$ by twisting or $\mathcal{U}_{h,g}(gl(2))$.

Let us take the particular nonlinear combinations of generators,

$$\begin{aligned}
A &= Z, \\
H' &= e^{-\sigma/2} H \\
X &= \frac{1}{2h} \sigma, \\
Y &= e^{-\sigma/2} \left(X_- + \frac{h}{2} H^2 \right) - \frac{h}{8} e^{\sigma/2} (e^{-\sigma} - 1).
\end{aligned} \tag{3.5}$$

Then A, H', X and Y satisfy the commutation relations,

$$\begin{aligned}
[X, Y] &= H', & [H', X] &= 2 \frac{\sinh hX}{h}, \\
[H', Y] &= -Y(\cosh hX) - (\cosh hX)Y, \\
[A, \bullet] &= 0.
\end{aligned} \tag{3.6}$$

The Hopf algebra mappings for these generators are given by

$$\begin{aligned}
\Delta(A) &= A \otimes 1 + 1 \otimes A, \\
\Delta(H') &= H' \otimes e^{hX} + e^{-hX} \otimes H' - \frac{2g}{h} \sinh hX \otimes Ae^{hX}, \\
\Delta(X) &= X \otimes 1 + 1 \otimes X, \\
\Delta(Y) &= Y \otimes e^{hX} + e^{-hX} \otimes Y - \frac{g^2}{h} \sinh hX \otimes A^2 e^{hX} + gH' \otimes Ae^{hX}, \\
\epsilon(\bullet) &= 0, \\
S(A) &= -A, & S(X) &= -X, \\
S(H') &= -e^{hX} H' e^{-hX} - \frac{2g}{h} (\sinh hX) A, \\
S(Y) &= -e^{hX} Y e^{-hX} + \frac{g^2}{h} (\sinh hX) A^2 + g e^{hX} H' A e^{-hX}.
\end{aligned}$$

Therefore the algebra generated by $\{A, H', X, Y\}$ is nothing but the one introduced in [4]. The authors of [4] define the algebra so as to be dual to the Jordanian matrix quantum group $GL_{h,g}(2)$ [5, 6].

4 Twist for $sl(1/2)$

The Lie superalgebra $sl(1/2)$ has four even and four odd elements denoted by $Z, H, X_{\pm}$ and $v_{\pm}, \bar{v}_{\pm}$, respectively. They satisfy the relations

$$[H, X_{\pm}] = \pm 2X_{\pm}, \quad [X_+, X_-] = H, \quad [Z, \bullet] = 0,$$

$$\begin{aligned}
[X_{\pm}, v_{\mp}] &= -v_{\pm}, & [X_{\pm}, \bar{v}_{\mp}] &= \bar{v}_{\pm}, \\
[Z, v_{\pm}] &= v_{\pm}, & [Z, \bar{v}_{\pm}] &= -\bar{v}_{\pm}, \\
[H, v_{\pm}] &= \pm v_{\pm}, & [H, \bar{v}_{\pm}] &= \pm \bar{v}_{\pm}, & [X_{\pm}, v_{\pm}] &= [X_{\pm}, \bar{v}_{\pm}] = 0, \\
\{v_{\pm}, v_{\pm}\} &= \{\bar{v}_{\pm}, \bar{v}_{\pm}\} = \{v_{\pm}, v_{\mp}\} = \{\bar{v}_{\pm}, \bar{v}_{\mp}\} = 0, \\
\{v_{\pm}, \bar{v}_{\pm}\} &= X_{\pm}, & \{\bar{v}_{\pm}, v_{\mp}\} &= \frac{1}{2}(Z \pm H).
\end{aligned} \tag{4.1}$$

It is easily seen from the above relations that the even elements $\{Z, H, X_{\pm}\}$ form a $gl(2)$ subalgebra and the universal enveloping algebra $\mathcal{U}(sl(1/2))$ is generated by $\{Z, H, X_{\pm}, v_{\pm}, \bar{v}_{\pm}\}$. The observation of $gl(2) \subset sl(1/2)$ implies that the twist element for $gl(2)$ can be used to twist $sl(1/2)$. Using (3.2), the twisted coproduct for the odd elements are given by

$$\begin{aligned}
\Delta(\bar{v}_{+}) &= \bar{v}_{+} \otimes e^{-\sigma/2} + \exp\left(-\frac{g}{2h}\sigma\right) \otimes \bar{v}_{+}, \\
\Delta(v_{-}) &= v_{-} \otimes e^{\sigma/2} + \exp\left(\frac{g}{2h}\sigma\right) \otimes v_{-} + hH \exp\left(\frac{g}{2h}\sigma\right) \otimes v_{+}e^{\sigma} \\
&\quad - gv_{+}e^{\sigma} \otimes Ze^{\sigma/2} - g(e^{\sigma} - 1) \exp\left(\frac{g}{2h}\sigma\right) \otimes Zv_{+}e^{\sigma},
\end{aligned} \tag{4.2}$$

and the antipode is

$$\begin{aligned}
S(\bar{v}_{+}) &= -\bar{v}_{+} \exp\left(\frac{1}{2h}(h+g)\sigma\right) \\
S(v_{-}) &= -(v_{-} - hHv_{+} + gv_{+}(1+Z)) \exp\left(-\frac{1}{2h}(h+g)\sigma\right).
\end{aligned} \tag{4.3}$$

The counit is undeformed and given by $\epsilon(\bar{v}_{+}) = \epsilon(v_{-}) = 0$. The Hopf algebra mappings for even elements have already been given in (3.3).

Definition 2. The triangular Hopf algebra generated by $\{Z, H, X_{\pm}, \bar{v}_{+}, v_{-}\}$ satisfying the relations (4.1), (4.2) and (4.3) is said to be the two-parametric deformation of $\mathcal{U}(sl(1/2))$ by twisting or $\mathcal{U}_{h,g}(sl(1/2))$.

The universal R-matrix for $\mathcal{U}_{h,g}(sl(1/2))$ is same as the one for $\mathcal{U}_{h,g}(gl(2))$. Noting that the fundamental representation of $\mathcal{U}_{h,g}(sl(1/2))$ is same as $sl(1/2)$, we obtain the R-matrix in the fundamental representation of $\mathcal{U}_{h,g}(sl(1/2))$, and see that it is a direct sum of four matrices

$$R = (1) \oplus \check{R} \oplus \check{R}^{-1} \oplus \bar{R}, \tag{4.4}$$

where

$$\check{R} = \begin{pmatrix} 1 & 2g \\ 0 & 1 \end{pmatrix}, \quad \bar{R} = \begin{pmatrix} 1 & h+g & -h-g & h^2-g^2 \\ 0 & 1 & 0 & h-g \\ 0 & 0 & 1 & -h+g \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{4.5}$$

The matrix $\bar{R}$ is the R-matrix (3.4) in the fundamental representation of $gl(2)$.

5 Two-parametric deformation of $SL(1/2)$

Using the R-matrix (4.4) and $\mathbf{Z}_2$ graded version of FRT-formalism [7], a matrix quantum supergroup dual to $\mathcal{U}_{h,g}(sl(1/2))$ can be constructed.

Introducing a 3×3 supermatrix

$$M = \begin{pmatrix} e & \Psi \\ \Theta & T \end{pmatrix},$$

where

$$\Psi = (\xi, \eta), \quad \Theta = \begin{pmatrix} \gamma \\ \delta \end{pmatrix}, \quad T = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

and e, T are even elements and Ψ, Θ are odd elements, the FRT-formalism guarantees that the commutation relations for the entries of M are given by RMM-relation and their Hopf algebra mappings are given by

$$\Delta(M) = M \otimes M, \quad \epsilon(M) = I_3, \quad S(M) = M^{-1}. \quad (5.1)$$

The commutation relations for the entries of M read

$$\begin{aligned} e\Psi &= \Psi e\check{R}, & \check{R}e\Theta &= \Theta e, & \check{R}eT &= Te\check{R}, \\ (\xi\Psi \ \eta\Psi) &= -(\Psi\xi \ \Psi\eta)\bar{R}, & \bar{R} \begin{pmatrix} \gamma\Theta \\ \delta\Theta \end{pmatrix} &= - \begin{pmatrix} \Theta\gamma \\ \Theta\delta \end{pmatrix}, \\ \check{R}(\xi\Theta \ \eta\Theta)\check{R} &= - \begin{pmatrix} \gamma\Psi \\ \delta\Psi \end{pmatrix}, & \bar{R} \begin{pmatrix} \gamma T \\ \delta T \end{pmatrix} &= \begin{pmatrix} T\gamma \\ T\delta \end{pmatrix} \check{R}, \\ \check{R}(\xi T \ \eta T) &= (T\xi \ T\eta)\bar{R}, & \bar{R}T_1T_2 &= T_2T_1\bar{R}. \end{aligned} \quad (5.2)$$

The last relation shows that the submatrix T satisfy the same algebraic relations as the $GL_{h,g}(2)$ [6]. As in [6], a determinant for T is defined by $\det T = ad - bc - (h+g)ac$, then it is not difficult to see that $\det T$ is not a center of deformed $SL(1/2)$ and the noncommutativity is independent of h

$$\begin{aligned} [\Omega, \det T] &= 0, \quad \Omega = e, \ \xi, \ \delta, \ c \\ [\eta, \det T] &= -2g\xi \det T, \quad [\gamma, \det T] = 2g\delta \det T, \\ [a, \det T] &= [\det T, d] = 2g c \det T, \\ [b, \det T] &= 2g\{(\det T)d - a(\det T)\}. \end{aligned}$$

Assuming that the $\det T$ is invertible, the explicit form of the inverse matrix of T can be obtain (see [6] for the formula).

We define a superdeterminant for the quantum supermatrix M by

$$s\det M = (\det T)^{-1}(e - \Psi T^{-1}\Theta). \quad (5.3)$$

This has the same form as the undeformed case (it is also called Berezinian in undeformed case). Direct computation shows that the $sdet M$ commute with all elements of the deformed $SL(1/2)$ so that we can safely set $sdet M = 1$. A coproduct and a counit for M are obvious from (5.1), however, an antipode is not. It is necessary to assume that the combination $e - \Psi T^{-1} \Theta$ has a inverse. Then the inverse matrix of M is given by [3]

$$M^{-1} = \begin{pmatrix} 1 & 0 \\ -T^{-1}\Theta & I_2 \end{pmatrix} \begin{pmatrix} (e - \Psi T^{-1}\Theta)^{-1} & 0 \\ 0 & T^{-1} \end{pmatrix} \begin{pmatrix} 1 & -\Psi T^{-1} \\ 0 & I_2 \end{pmatrix}, \quad (5.4)$$

where I_2 is the 2×2 unit matrix.

Definition 3. An algebra generated by the entries of M satisfying (5.2), (5.1), (5.4) and $sdet M = 1$ is said to be the two-parametric deformation of $Fun(SL(1/2))$ or $SL_{h,g}(1/2)$.

We finally give some remarks. The twist element $\mathcal{F}$ (3.2) is also applicable to deform $osp(1/2) \oplus u(1)$, since $gl(2) \subset osp(1/2) \oplus u(1)$. The universal R-matrix for the obtained algebra is used to deform the supergroup $OSp(1/2) \otimes U(1)$. The inclusion of a odd element of $sl(1/2)$ into a twist element may be possible. The inclusion of a odd elements is found for $osp(1/2)$ recently [14].

References

- [1] Ballesteros A., Herranz J., and Parashar P.: Multiparametric quantum $gl(2)$: Lie bialgebras, quantum R -matrices and non-relativistic limits, Preprint math.QA/9806149.
- [2] Drinfeld V. G.: Leningrad Math. J. **1** (1990) 1419.
- [3] Celeghini E. and Kulish P. P.: J. Phys. A: Math. Gen. **31** (1998) L79.
- [4] Aneva B. L., Dobrev V. K., and Mihov S. G.: J. Phys. A: Math. Gen. **30** (1997) 6769.
- [5] Demidov E. E. *et al*: Progr. Theor. Phys. Suppl. **102** (1990) 203.
- [6] Aghamohammadi A.: Mod. Phys. Lett. A **8** (1993) 2607.
- [7] Faddeev L. D., Reshetikhin N. Yu., and Takhtajan L. A.: Leningrad Math. J. **1** (1990) 193.
- [8] Reshetikhin N.: Lett. Math. Phys. **20** (1990) 331.

- [9] Ohn Ch.: Lett. Math. Phys. **25** (1992) 85.
- [10] Lukierski J., Ruegg H., and Tolstoy V. N.: J. Phys. A:Math. Gen. **27** (1994) 2389.
- [11] Bonechi, F. **et al**: Comm. Math. Phys. **169** (1995) 627, Abdesselam, B.: J. Math. Phys. **38** (1997) 6045.
- [12] Kulish P. P., and Mudrov A. I.: Universal R -matrix for esoteric quantum group, Preprint math.QA/9804006.
- [13] Kulish P. P, Lyakhovsky V. D., and Mudrov A. I.: Extended jordanian twists for Lie algebras, Preprint math.QA/9806014.
- [14] Kulish P. P: Super-jordanian deformation of the orthosymplectic Lie superalgebra, Preprint math.QA/9806104.